



Forensic Medicine Authority
Egyptian Journal of Forensic Sciences

www.sciencedirect.com



Suicide rate: Trends and implications in Upper Egypt

Wafaa M. Abdel Moneim, Heba A. Yassa *, Saffa M. George

Forensic Medicine and Clinical Toxicology Department, Faculty of Medicine, Assiut University, Egypt

KEYWORDS

Suicide;
 Assiut;
 Province

Abstract Suicide is an important problem, ranking among the top 10 causes of death for individuals in all ages in developed countries. The present study is a retrospective study evaluating suicide cases in Assiut, one of the largest province in Egypt from 2005 to 2009. There were 117 cases of which involved 68 male victims (58.11%) and involved 49 females (48.11%). Suicide rates ranged from (0.6 to 0.8 per 100,000). Age predominance was 20–≤30 years. The method of suicide was different between male and female victims, as male victims tried to use more violent methods than females. *Conclusion:* the present study showed that suicide rates have increased since 1987, indicating a grave problem that needs to be solved.

© 2011 Forensic Medicine Authority. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

Suicide is an important problem, ranking among the top 10 causes of death for individuals of all ages in developed countries. It is believed that the most dramatic increase in suicide mortality will be observed in third world countries because of socioeconomic and behavioral factors.¹ Every year, almost one million people die as a result of suicide, with a mortality rate of about 16 per 100,000, which means that one death occurs every 40 s. In the last 45 years, suicide rates worldwide have increased by 60%. Suicide is among the three leading causes of death among people aged 15–44 years in some coun-

tries and the second-leading cause of death among those aged 10–24 years; these figures do not include the suicide attempts, which are up to 20 times more frequent than completed suicide.²

The most important problem in Egypt is that there is no data bank for suicide cases. This causes the suicide problem to be underestimated and, thus, neglected by the government, although this problem seems to be universal.^{3–6} The last data found regarding suicide rates in Egypt, according to the WHO, was in 1987.⁷

According to the WHO's regional distribution, the lowest rate of suicide was found in the Eastern Mediterranean region, which comprises mostly Arabic countries, but this researcher feels that this is due to the underestimation of suicide cases in this area. In the Eastern Mediterranean region (EMR), suicide death rates are still low in comparison to those of the European, Pacific, and Asian regions. In 2000, suicide was estimated to be the 25th leading cause of death in the Eastern Mediterranean region (EMR) but was ranked 7th in the European region, 8th in the Western Pacific region, and 16th in the South East Asian region.⁸

A great reason that Egypt has received little attention from suicidologists is that suicide has not been looked upon as a problem of any importance; there is a wish to deny its existence. This attitude may be explained partly by the fact that

* Corresponding author. Tel.: +20 101349209.

E-mail address: heba_yassa@hotmail.com (H.A. Yassa).



Table 1 Distribution of suicidal cases according to age group per year.

Year		Suicidal frequency per age group																% Suicid rate per 100,000	
		0–≤10		10–≤20		20–≤30		30–≤40		40–≤50		50–≤60		60–≤70		70–≤80		Total	%
2005	M	0	0	3	5	6	9	3	7	1	2	2	2	0	1	1	1	16	26
	F	0	0	2	3	4	5	1	2	0	0	0	0	0	0	0	0	13.67	22.2
2006	M	0	0	0	2	0	5	2	5	1	2	2	2	1	2	2	2	8	20
	F	0	0	2	5	3	1	0	0	0	0	0	0	0	0	0	0	6.83	17.1
2007	M	0	0	0	2	8	15	4	6	1	2	1	2	1	1	0	0	15	28
	F	0	0	2	7	2	1	1	1	0	0	0	0	0	0	0	0	12.82	23.9
2008	M	0	0	1	4	2	5	3	3	3	3	2	2	1	1	0	1	12	19
	F	0	0	3	3	0	0	0	0	0	0	0	0	0	1	0	0	10.25	16.2
2009	M	0	0	1	4	7	9	4	6	3	3	1	1	1	1	0	0	17	24
	F	0	0	3	2	2	0	0	0	0	0	0	0	0	0	0	0	14.53	20.5
Total		0	0	17	43	27	12	9	5	4	117	100							

Egyptians, from the dawn of their history, have worshipped life.⁹ Religion also may play a great role in giving people defenses against isolation or alienation.¹⁰ There is a general agreement that previous statistics on the incidence of suicide in Egypt are grossly inadequate and that their comparison based on the figures available is inaccurate and misleading.⁹ In addition, the features of deaths due to suicide are changing constantly, revealing complex social and cultural developments.¹¹ This idea supported by Rezaeian (2010),²² who highlighted that, within Muslim-dominated Middle Eastern countries, suicide rates are increasing among young females, and the study confirmed that there are high rates of suicide and attempted suicide among young Muslim females in the Middle East. Furthermore, Prichard and Amanullah (2007)²³ suggested that, to avoid the under-reporting of suicides, both formal suicide verdicts and other violent deaths should be considered together because the other violent deaths may include hidden suicides.

1.1. Aim of the study

- (1) To investigate different causes of suicide among those living in Upper Egypt.
- (2) To compare the suicide rates among different age groups and between sexes.
- (3) To compare the method of suicide between male and female victims.

2. Type of study: retrospective study

2.1. Subjects and method

2.1.1. Data collection

Data on suicide incidence, age, sex, residence (urban or rural), season, previous attempts, presence or absence of mental disease history, and methods of suicides were collected retrospectively from the Forensic Medicine Institute in the Assiut province from 2005 to 2009. The Forensic Medicine Institute reports include full autopsy and toxicological analysis for the cause of death to differentiate between cases of homicidal, suicidal, or accidental cause of death.

The Assiut province (one of the biggest Upper Egypt provinces) has a population of about 3444,967, according to 2006 statistics.²⁰ The Assiut province is a large one

Containing 10 cities: Dairout, El Kousia, Abnoub, Manfalout, Assiut, El Fath, Abouteeg, Elghanaym, Sahel Seleem, El Badary, and Sedfa.¹²

Suicide methods were classified into seven groups according to the cause of death: poisoning by drugs, poisoning by other means, hanging, drowning, firearm, falling from a height, and burns.

2.2. Statistical analysis

The suicide rates in the Assiut province were calculated for each year from 2005 to 2009 and classified according to sex and age group. In addition, the youth suicide rates in Assiut were compared to the elderly suicide rate, and the rates between males and females were compared. The comparison between males and females was done to assess suicidal tendencies, and the differences between them with respect to pattern of death were evaluated with a confidence interval of 95% using SPSS program version 15.

3. Results

The suicide data collected in the period from 2005 to 2009 is shown distributed by year and age group in Table 1. This table shows that the total number of cases of suicide in Assiut during this period was 117. The suicide rate per 100,000 ranged from 0.551 in 2008 to 0.813 in 2007. Table 2 shows the distribution of suicide cases among males and females per year. It shows that the rate for males was 58.11% and the rate for females was 41.89%. The suicide cases were concentrated in the age group of 20–≤30 years (36.7%), followed by the age group of 30–≤40 years (10.25%), which is the full youth age range,

Table 2 Distribution of suicidal cases according to the sex.

Year	Male	Female	Total
2005	16	10	26
2006	8	12	20
2007	15	13	28
2008	12	7	19
2009	17	7	24
Total	68	49	117
%	58.11	41.88	100




Figure 1 Distribution percent of suicidal cases per year in Upper Egypt (2005–2009).




Figure 2 Distribution of suicidal deaths per year and age group.

and this indicates a big problem. The table also shows that the tendency of males to commit suicide was more than that of females in the same age groups; this is shown in Figs. 1 and 2.

Fig. 3 shows the distribution of cause of death in suicide cases among males. It shows that suicide using toxins was the most common method among males (29%), followed by hanging (28%), firearm (16%), burns (12%), drowning (10%), and, finally, falling from a height (5%). Furthermore, Fig. 4 shows the distribution of cause of death in suicide cases among females. It shows that suicide by toxins is the first most common cause of suicide among females (70%), followed by burns (12%), drowning and falling from a height (6% each), hanging (4%), and, finally, firearm (2%).

Table 3 and Fig. 5 show the most common toxins used in suicide among males and females. It shows that the number of cases among females is greater than that of males.

The most common toxin used among both males and females was organophosphates. Other toxins used were, carbamate, lanet, phenols, phosphorous, strychnine, and toxafen.

Fig. 6 shows the distribution of the site of firearm injuries in suicide cases among males and females. It shows that the most common area for firearm injury was the forehead (42%), followed by the abdomen (25%), chest (17%), and roof of the mouth and neck (8% each).




Figure 3 Cause of death in male suicidal cases in the period of (2005–2009).




Figure 4 Cause of death in female suicidal cases in the period of (2005–2009).

Table 3 Most common toxins used in suicide among male and females.

Type of toxins	Sex		Total	Percent
	Male	Female		
Drug overdose	0	1	1	1.85
Organophosphate	13	28	41	75.9
Carbamate	1	1	2	3.7
Lanet	3	1	4	7.4
Phenols	1	0	1	1.85
Phosphorous	1	1	2	3.7
Strychnine	1	0	1	1.85
Toxafen	1	1	2	3.7
Total	21	33	54	100




Figure 5 Distribution percent of toxins used in suicide among male and female in Upper Egypt.




Figure 6 Distribution percent of site of the firearm injuries used in suicide among male and female in Upper Egypt.

4. Discussion

The present study evaluated the rates, trends, and distribution of suicide from 2005 to 2009 in Assiut, one of the largest cities in Upper Egypt. The average rate of suicide during the study period ranged from 0.6 to 0.8 per 100,000 persons. The rate substantially exceeded the 0.1 and 0.0 (male and female, respectively) per 100,000 persons reported by the WHO for Egypt in 1987. This may be explained by the effect of increasing depression among people, which may be aggravated due to social problems such as unemployment and spinsterhood. An additional factor that may aggravate the state of depression in youth, especially among males in our area, is the increase in drug abuse among youth, which is difficult to explain: does the depression lead them to use these drugs or does the drug abuse increase depression? This idea is supported by the work of Yassa et al.²⁵ who indicated that bango (cannabis leaves), commonly abused drug, was widely spread among youth in Upper Egypt, concentrated in the age group between 21 and ≤ 31 years, which is the same age group that has the highest suicide rate.

The fact that the suicide rate is still lower than that of other countries may be explained by the effect of religion and the good relations between family members throughout the country, especially in Upper Egypt. This idea is supported by the work of Setenay et al.²⁴ who reported that religion plays a great role, especially with respect to a belief in an afterlife, which averts feelings of hopelessness, a feeling that has been described as an important predisposing factor for suicide.

The present study showed that there were gender differences in suicide rates, with rates for male rate being higher for most of the period of the study. This may be explained by the increasing stress on males, mainly in Upper Egypt, where the work load is greater for them. These results may be compared with those of other countries, as in the work of Bjerkeset et al. (2008), who demonstrated in a follow-up study of failed suicide attempts over a period of 8 years that there was a higher suicide risk in men than in women, as they are more susceptible to anxiety and depression.^{5,13–15}

The age distribution for males and females appeared to be in the age range of 20– ≤ 30 years and the age group of 30– ≤ 40 years, which is the age of youth and work and, in females, the age of reproduction. This indicates the existence of a problem: a decrease in the chance of working and the chance of marriage in females. Gad ElHak (2009) supported these results in demonstrating that the age group with the highest rate of suicide in Port Said, another city in Egypt, was in the age

group of 20–30 years. The suicide rate decreased as age increased among males and females, which can be explained by the increase in the level of responsibility of males and females toward their families.

The causes of death among males differed from those of females, as shown in Figs. 3 and 4. In males, the main cause of death was toxins (29%), followed by hanging (28%), firearm (16%), burns (12%), drowning (10%), and falling from a height (5%), while, in females, the distribution of causes of death was as follows: toxins (70%), followed by burns (12%), drowning (6%), falling from a height (6%), and firearm injuries (2%). This can be explained by the differences in the personality characteristics between males and females, as males mostly chose more violent methods of suicide. These results matched those of Hawton (2000), who stated that the suicide rates in most countries are higher among males than in females, except for China, which has very high rates of suicide in females, especially in rural areas. This is explained by the fact that females seek help for psychological problems more than males.^{15–17} These results differ from those of the WHO record, which classify the methods of suicide in different countries according to the WHO mortality database, finding that, in most of the studied countries (not including Egypt), hanging was the most frequent cause of suicide among males, followed by firearm injuries and poisoning.²¹

In the present study, the most common toxin used for suicide among males and females was organophosphates, comprising 75.9% of all suicidal cases, followed by lanet, carbamate, and toxafen (7.4%, 3.7%, and 3.7%, respectively). Jaga and Dharmani (2007)¹⁸ found that there was a relation among populations exposed to organophosphates, which leads to neurobehavioral effects, depression, suicide, and death. Furthermore, London et al. (2005)¹⁹ found that suicide rates were high in farming populations. They found that exposure to organophosphates led to serotonin disturbances, which are implicated in depression and suicide.

In the present study, suicide by firearm injuries was more common for males, and the most common area found to be affected was the forehead (42%), followed by the abdomen (25%), chest (17%), neck (8%), and roof of the mouth (8%).

5. Conclusion

Suicide is a very grave problem that may be caused and aggravated by many factors that occur in our society, such as unemployment and spinsterhood. Suicide affects mainly youth who are capable of work, which causes a big loss to the community. Therefore, all efforts should be directed to solving youths' problems to decrease the rate of suicide in our society.

Acknowledgements

We thank Dr. Abdel Karim Hassan of the Assiut Forensic Medicine Institute, who facilitates our mission at the institute.

References

- Gad ElHak S, El-Ghazali A, Salama M, Aboelyazeed A. Fatal suicide cases in Port Said city, Egypt. *J Foren Legal Med* 2009;16:168–266.

2. World Health organization report, "Suicide Prevention". <http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/>f accessed in 13/8/2010 (2009).
3. Barraclough B. Differences between national suicide rates. *Br J Psychiatry* 1973;**122**:95–6.
4. Sainsbury P, Jenkins J. The accuracy of officially reported suicide statistics for purposes of epidemiological research. *J Epidemiol Commun Heal* 1982;**36**:43–8.
5. Bertolote J, Fleischmann A. A global perspective in the epidemiology of suicide. *Suicidology arg* 2002;**7**(2):6–8.
6. Chang S, Sterene J, Lu T, Gunnell D. Hidden suicides amongst deaths certified as underdetermined intent accident by pesticide poisoning and accident by suffocation in Taiwan. *Soc Psychiatry Psychiatr Epidemiol* 2009, doi: 10.1007.
7. World Health Organization report: WHO report suicidal statistics 2011. Who.int. 2011-01-10. Retrieved 2011-01-10.
8. Rezaeian M. Age and sex suicide rates in the Eastern Mediterranean region based on global burden of disease estimate for 2000. *East Mediterr Health J* 2007;**13**(4):953–60.
9. Okasha A, Lotaeif F. Suicide in Egypt. 1st ed. In: Headdley L, editor. *Suicide in Asia and Near East*. London: England press; 1983. p. 333–50 [Chapter 14].
10. Stengel E. Suicide in Egypt. 1st ed. In: Headdley L, editor. *Suicide in Asia and Near East*. London: England press; 1969. p. 333–50 [Chapter 14].
11. Byard R, Kliitte A, Gilbert J. Changing patterns of female suicide: 1986–2000. *J Clin Forensic Med* 2004;**11**:123–8.
12. www.wikipedia.com Assiut, accessed in 17/9/2010.
13. Rohling J, Friend J, Powell A. Adolescent suicide, gender, and culture: a rate and risk factor analysis. *Aggress violent Behav* 2009;**14**:402–14.
14. Inoue K, Tanii H, Kaiya H, Abe S, Nishimura Y, Masaki M, Okazaki Y, Nata M, Fukunaga T. The correlation between unemployment and suicide rates in Japan between 1978 and 2004. *Leg Med* 2007;**9**:139–42.
15. Bjerkeset O, Romundstad P, Gunnell D. Gender differences in the association of mixed anxiety and depression with suicide. *Br J Psychiatry* 2008;**192**:474–5.
16. Varnik A, Kolves K, Van Der Feltz-Cornelis C, Marusic A, Oskarsson H, Palmer A, Reisch T, Scheerder G, Arensman E, Giupponi G, Gusmao R, Maxwell M, Pull C, Szekely A, Sola VP, Hegerl U. Suicide methods in Europe—a gender specific analyses of countries participating in the European Alliance Against Depression. *J Epidemiol Commun Health* 2008;**62**:545–51.
17. Hawton K. Sex and suicide, gender differences in suicidal behavior. *Br J Psychiatry* 2000;**177**:484–5.
18. Jaga K, Dharmani C. The interrelation between organophosphates toxicity and the epidemiology of depression and suicide. *Rev Env Health* 2007;**22**(1):57–73.
19. London L, Flisher A, Wesseling C, Mergler D, Kromhout H. Suicide and exposure to organophosphates insecticides: cause or effect? *Am J Ind Med* 2005;**47**(4):308–21.
20. Central Agency for Public Mobilization and Statistics Egypt, City population. In: <<http://www.citypopulation.de/Egypt.html>> accessed in 14/9/2010 (2006).
21. Ajdaic-Gross V, Weiss M, Ring M, Hepp U, Bopp M, Gutzwiller W, Rossler W. Methods of suicide: international suicide patterns derived from the WHO mortality database. *Bull World Health Organ* 2008;**86**(9):726–32.
22. Rezaeian M. Suicide among young Middle Eastern females. *Crisis* 2010;**31**(1):36–42.
23. Pritchard C, Amanullah S. An analysis of suicide and underdetermined deaths in 17 predominantly Islamic countries contrasted with the UK. *Psychol Med* 2007;**37**(3):421–30.
24. Setenay O, Yenilmez C, Ayranci U, Gunay Y, Ozdamar K. Sexual differences in the completed suicides in Turkey. *Eur Psychiatry* 2007;**22**(4):223–8.
25. Yassa H, Dawood A, Shehata M, Abdel Hady R, Abdel Aal K. Risk factors for bango abuse in Upper Egypt. *Environ Toxicol Pharmacol* 2009;**28**:397–402.